



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CORRESPONDENCE.

ON THE METHOD OF DERIVING FROM A TABLE OF PREMIUMS
THE RATE OF MORTALITY ON WHICH IT IS BASED.*To the Editor of the Assurance Magazine.*

SIR,—I have looked over several prospectuses of Assurance Companies lately, and have been puzzled in some cases to find out the table of mortality according to which the annual premiums have been calculated throughout. Either the pure premiums have been loaded capriciously, or the premium tables have been meddled with by unskilful hands, in some instances; or, possibly, many Offices have their own particular tables of mortality, which are not made known to the world; or perhaps there may be some whose dexterity has enabled them to graduate a table of premiums irrespective of any table of mortality whatever.

I shall therefore assume that a table of pure premiums is given, to find out the table of mortality upon which it is based.

Let p_m and p_{m+1} be the pure premiums for any two consecutive ages; a_m, a_{m+1} , the present values of annuities corresponding to these ages; l_m, l_{m+1} , the numbers living, and which are to be registered in the table

sought; $v = \frac{1}{1+r}$ and $d = 1 - v$. Then (1), $p_m = \frac{1}{1+a_m} - d$; (2), $p_{m+1} = \frac{1}{1+a_{m+1}} - d$; also (3), $a_m = \frac{l_{m+1}v}{l_m}(1+a_{m+1})$. By means of these three equations the values a_m and a_{m+1} may be eliminated thus—

$$p_{m+1} - p_m = \frac{1}{1+a_{m+1}} - \frac{1}{1+a_m} \text{ from (3) } \frac{1}{1+a_{m+1}} = \frac{l_{m+1}v}{l_m a_m}$$

$$\therefore p_{m+1} = \frac{l_{m+1}v}{l_m a_m} - d, \text{ and } a_m = \frac{v - p_m}{d + p_m}.$$

Hence

$$p_{m+1} + d = \frac{l_{m+1}v}{l_m} \cdot \frac{d + p_m}{v - p_m}, \text{ or } \frac{(p_{m+1} + d)(v - p_m)l_m}{v(d + p_m)} = l_{m+1},$$

in which l_m may be assumed arbitrarily. Suppose $l_m = l_{10}$, then l_{10} being made equal to any convenient number, l_{11}, l_{12}, l_{13} , &c. may be found throughout.

If $l_{10} = 6460$, then $p_{10} = \cdot 0116702$; $p_{11} = \cdot 0119812$; $d = \cdot 0291260$;
 $v = \cdot 9708740$ $\therefore l_{11} = \frac{(\cdot 0411072)(\cdot 9592038)(6460)}{(\cdot 0407962)(\cdot 970874)} = 6431$, as in the Carlisle Table.

GEO. SCOTT.

40, King William Street, London Bridge,
 14th April, 1856.

NOTE.—A communication from Mr. William Wylie on this subject will be found at page 391, vol. ii., of this *Journal*.—ED. A. M.